

Citation for published version:

Gonzalez, J & Betts, J 2019, 'Dietary Sugars, Exercise and Hepatic Carbohydrate Metabolism: Sugars and liver metabolism', *Proceedings of the Nutrition Society*, vol. 78, no. 2, pp. 246-256.
<https://doi.org/10.1017/S0029665118002604>

DOI:

[10.1017/S0029665118002604](https://doi.org/10.1017/S0029665118002604)

Publication date:

2019

Document Version

Peer reviewed version

[Link to publication](#)

This article has been published in *Proceedings of the Nutrition Society* on 23 October 2018. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. (C) Cambridge University Press, 2018.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title: Dietary Sugars, Exercise and Hepatic Carbohydrate Metabolism

Authors: Javier T. Gonzalez and James A. Betts

Affiliation: Department for Health, University of Bath, Bath, BA2 7AY, United Kingdom.

Correspondence: Javier T. Gonzalez, Department for Health, University of Bath, Bath, BA2 7AY, United Kingdom.

E-mail: J.T.Gonzalez@bath.ac.uk Telephone: 0(+44)1225 385518

Short title: Sugars and liver metabolism

Keywords: Glucose; Galactose; Fructose; Glycogen; Physical activity

1 **Abstract**

2 This paper reviews the physiological responses of human liver carbohydrate metabolism to physical
3 activity and ingestion of dietary sugars. The liver represents a central link in human carbohydrate
4 metabolism and a mechanistic crux point for the effects of dietary sugars on athletic performance and
5 metabolic health. As a corollary, knowledge regarding physiological responses to sugar ingestion has
6 potential application to either improve endurance performance in athletes, or target metabolic
7 diseases in people who are overweight, obese and/or sedentary. For example, exercise increases
8 whole-body glycogen utilisation, and the breakdown of liver glycogen to maintain blood glucose
9 concentrations becomes increasingly important as exercise intensity increases. Accordingly,
10 prolonged exercise at moderate-to-high exercise intensity results in depletion of liver glycogen stores
11 unless carbohydrate is ingested during exercise. The exercise-induced glycogen deficit can increase
12 insulin sensitivity and blood glucose control, and may result in less hepatic lipid synthesis. Therefore,
13 the induction and maintenance of a glycogen deficit with exercise could be a specific target to
14 improve metabolic health and could be achieved by carbohydrate (sugar) restriction before, during
15 and/or after exercise. Conversely, for athletes, maintaining and restoring these glycogen stores is a
16 priority when competing in events requiring repeated exertion with limited recovery. With this in
17 mind, evidence consistently demonstrates that fructose-containing sugars accelerate post-exercise
18 liver glycogen repletion and could reduce recovery time by as much as half that seen with ingestion
19 of glucose (polymers)-only. Therefore, athletes aiming for rapid recovery in multi-stage events should
20 consider ingesting fructose-containing sugars to accelerate recovery.

21
22
23
24
25

26 **Introduction**

27 The French physiologist Claude Bernard (1813-1878) is not only one of the first to propose blind
28 experiments to reduce bias⁽¹⁾ but also is credited with the discovery of glycogen in the liver, thus
29 revealing the central role of this organ in the homeostatic regulation of blood glucose concentrations
30 (or “*milieu intérieur*”)^(2; 3). Bernard originally intended to study the metabolism of all types of foods,
31 choosing to start with the putatively *simple* metabolism of sugars. The complexities of sugar
32 metabolism led Bernard to focus on this area for more than 30 years and, in understated fashion, he
33 described this systematic and meticulous undertaking as “research which has not been wholly
34 sterile”⁽⁴⁾. During this time, he found that the portal vein of dogs (the major blood supply to the liver)
35 has little to no glucose, whereas the hepatic vein leaving the liver carries substantial quantities of
36 glucose. This led Bernard to conclude that the liver is a potential source of sugar. This capacity of the
37 liver to supply glucose to the systemic circulation is important when dietary carbohydrate intake is
38 insufficient to meet the carbohydrate demands of tissues such as the brain and muscles. Therefore,
39 during fasting, exercise or consumption of low-carbohydrate diets, the liver can supply glucose for
40 peripheral tissues. Glucose produced by the liver is derived from two sources: the breakdown of
41 stored glycogen (i.e. glycogenolysis), and the *de novo* production of new glucose from precursors
42 such as lactate, glycerol, pyruvate, glucogenic amino acids, fructose and galactose (i.e.
43 gluconeogenesis)⁽⁵⁾. The liver is the largest glycogen store in humans that can be hydrolysed and
44 release glucose into the circulation to sustain blood glucose concentrations, and is also the tissue with
45 the greatest capacity for gluconeogenesis. Therefore, the ability for the liver to supply glucose from
46 both glycogenolysis and gluconeogenesis has important consequences for maintaining metabolic
47 control during exercise, and especially when dietary carbohydrate intake is restricted.

48 The liver also plays a central role in the postprandial metabolism of carbohydrates. Following
49 intestinal absorption, the liver is one of the first tissues exposed to ingested carbohydrate. Whilst the
50 intestine (and kidneys) can also metabolise some dietary sugar⁽⁶⁾ and undertake gluconeogenesis⁽⁷⁾,
51 these are quantitatively less important than hepatic metabolism, at least in humans⁽⁸⁾. Various types
52 of sugars are distinctly metabolised by the liver, with potential implications for human health and
53 performance^(9; 10). Accordingly, the aim of this narrative review is to describe the hepatic metabolism
54 of dietary sugars at rest and during exercise, while considering potential implications for human
55 health and (endurance) exercise performance.

56

57 **Dietary Sugars**

58 Common sugars in the human diet include the monosaccharides: glucose, fructose and galactose; and
59 the disaccharides: sucrose (fructose-glucose), lactose (galactose-glucose) and maltose (glucose-
60 glucose)⁽⁹⁾. Dietary sugars can be consumed from a variety of food sources, which can influence

61 resultant health effects. The World Health Organization (WHO) classifies sugars into those which
62 are “intrinsic” (e.g. incorporated within the structure of intact fruit and vegetables or lactose/galactose
63 from milk) *versus* “free sugars”⁽¹¹⁾. Free sugars are defined by WHO as monosaccharides and
64 disaccharides added to foods and beverages by the manufacturer, cook or consumer, along with sugars
65 naturally present in honey, syrups, fruit juices and fruit juice concentrates. This classification system
66 is useful for distinguishing between food sources of sugar that are energy dense (i.e. free sugars) and
67 thus may contribute to weight gain^(11; 12). However, this classification system does not specifically
68 distinguish between ingestion of glucose-containing sugars and fructose- or galactose-containing
69 sugars in relation to health, nor does it consider the physical activity status of the individual. This is
70 interesting considering the fundamental differences in the intestinal absorption and hepatic
71 metabolism of glucose, fructose and galactose, and how this metabolism is altered during exercise^{(8;}
72 9; 10).

73 Before describing the hepatic metabolism of carbohydrates and sugars, it is important to
74 clarify two points. First, the hydrolysis of glucose polymers such as maltodextrin and starch are
75 typically not rate limiting to intestinal absorption⁽¹³⁾, and therefore (at least with regards to hepatic
76 metabolism) free glucose, maltose, maltodextrin and starch can all be considered physiologically
77 similar stimuli. Second, in typical human diets, free glucose is rarely consumed alone, but rather is
78 usually consumed alongside fructose or lactose, or is consumed in polymer (non-sugar) form, such
79 as maltodextrin and starch. Accordingly, whilst this review will refer to the specific types of sugars
80 utilised in studies (e.g. glucose only vs fructose-glucose mixtures), it can be viewed that a non-
81 fructose or non-galactose condition (such as glucose or maltodextrin ingestion) is physiologically
82 representative of non-sugar intake (i.e. maltodextrin, starch etc.), whereas fructose-glucose and
83 galactose-glucose co-ingestion represent the physiological responses to typical sugar intake.


84 **Hepatic Metabolism of Sugars at Rest**

85 Glucose and galactose are primarily absorbed across the intestinal lumen via the transport
86 protein sodium-dependent glucose transporter 2 (SGLT2)⁽¹⁴⁾, whereas fructose is primarily absorbed
87 via glucose transporter 5 (GLUT5)⁽¹⁴⁾. Once absorbed, these sugars are then metabolised very
88 differently. Glucose is preferentially metabolised by extra-splanchnic tissues such as skeletal muscle,
89 the brain and cardiac muscle^(15; 16), whilst fructose and galactose are primarily metabolised by the
90 liver and to a lesser extent small bowel enterocytes and proximal renal tubules^(10; 17). Peripheral tissues
91 such as muscle are therefore only exposed to relatively small amounts of fructose and galactose^{(18;}
92 ¹⁹⁾.

93 Compared to galactose, the metabolic fate of fructose is relatively well characterised. At rest,
94 the liver rapidly takes up and metabolises fructose into fructose-1-phosphate (P) via fructokinase (K_m
95 for fructose: ~ 0.5 mM and V_{max} estimated at ~ 3 mmol \cdot min⁻¹ \cdot g human liver⁻¹)^(20; 21; 22). Fructose-1-P is
96 then metabolised into triose-P (3 carbon substrates) via aldolase B⁽²³⁾. At rest, the majority of fructose-
97 derived triose-P is converted via gluconeogenesis into glucose ($\sim 50\%$) and glycogen (~ 15 - 25%), but
98 some of triose-P can be metabolised into pyruvate, then either oxidised within the liver or converted
99 into lactate ($\sim 25\%$), which enters the systemic circulation and can increase blood lactate
100 concentrations^(10; 24). One other fate of ingested fructose is the conversion into fatty acids via the
101 process known as *de novo* lipogenesis⁽²⁵⁾. It has been suggested that lactate is the primary precursor
102 to hepatic *de novo* lipogenesis with fructose intake⁽²⁶⁾, but the proportion of fructose that is ultimately
103 converted into lipid is estimated at $<1\%$ and therefore represents a quantitatively minor pathway of
104 disposal⁽¹⁹⁾. Nevertheless, the effects of ingested fructose on *de novo* lipogenesis may still be
105 important for metabolic health⁽²⁷⁾.

106 Quantitative estimates of the metabolic fate of galactose in humans are scarce. It has been
107 suggested that the primary pathway for human galactose metabolism is the Leloir pathway, the
108 enzymes of which show highest activity in the liver⁽¹⁷⁾. This pathway involves four main steps: 1)
109 phosphorylation of galactose by galactokinase (K_m for galactose: ~ 0.9 mM and V_{max} estimated at ~ 1.4
110 mmol \cdot min⁻¹ \cdot g human liver⁻¹ (28; 29)) to yield galactose-1-phosphate; 2) conversion of galactose-1-
111 phosphate and uridine diphosphate (UDP) glucose to UDP galactose and glucose-1-phosphate by
112 galactose-1-phosphate uridylyltransferase; 3) conversion of UDP-galactose to UDP-glucose by UDP-
113 galactose-4-epimerase; and 4) conversion of UDP-glucose and diphosphate (PPi) to glucose-1-
114 phosphate and uridine triphosphate (UTP) by UDP glucose pyrophosphorylase⁽¹⁷⁾. Of note, is an
115 alternative pathway for step 2, known as the Isselbacher pathway, whereby galactose-1-phosphate
116 and UTP are converted to UDP-galactose and PPi by the enzymes UDP-galactose pyrophosphorylase
117 and UDP-glucose pyrophosphorylase⁽¹⁷⁾. Some tracer studies have attempted to determine the
118 metabolic fate of oral galactose in humans, estimating that during ingestion of galactose at a rate of

119 33 $\mu\text{mol}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ (~ 135 g over 360 min), the splanchnic uptake of galactose is saturable at ~ 15
 120 $\mu\text{mol}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ^(30; 31). At this rate of ingestion, it is estimated that ~ 30 -60% of the ingested galactose
 121 is converted into glucose⁽³⁰⁾, mostly via the direct conversation of hexose to glucose ($\sim 67\%$), with
 122 some converted via the indirect (hexose to 3 carbon substrates to glucose) pathway ($\sim 33\%$)⁽³¹⁾.
 123 Ultimately, the metabolic fate of ingested galactose in humans therefore remains incompletely
 124 understood, although it has been speculated that liver glycogen synthesis is a major route^(32; 33)
 125 (Figure 1).



138 **Figure 1.** Major metabolic pathways of glucose, fructose and galactose in the human liver. TCA,
 139 tricarboxylic acid cycle. Based on references: 10, 17, 23, 24, 25, 26, 30, 31, 32, 33.

140 **Hepatic Metabolism of Carbohydrates with Exercise**

141 Exercise increases energy expenditure, which is predominantly met during prolonged (>30 min)
 142 exercise by increases in both carbohydrate and fat oxidation compared to the resting state⁽³⁴⁾. The
 143 relative contributions of carbohydrate versus fat to exercise metabolism are influenced by the
 144 intensity and mode of exercise⁽³⁵⁾, preceding nutritional status^(36; 37; 38), endurance training status⁽³⁹⁾
 145 and biological sex⁽⁴⁰⁾. Specifically, higher carbohydrate oxidation rates are seen with cycling versus
 146 running⁽³⁵⁾, higher versus lower exercise intensity⁽³⁴⁾, prior carbohydrate feeding versus fasting⁽³⁷⁾, in
 147 individuals who are less versus more endurance-trained⁽³⁹⁾ and amongst men *versus* women⁽⁴⁰⁾. Of
 148 these predictive factors, the intensity of exercise seems to be the most potent in determining
 149 carbohydrate and fat utilisation^(34; 41). Even in highly-trained athletes studied in the overnight fasted
 150 state, carbohydrates are the predominant fuel source during moderate-to-high intensity ($>50\%$
 151 $\dot{V}\text{O}_{2\text{peak}}$) exercise⁽³⁴⁾. Exercise is therefore a potent modulator of carbohydrate metabolism, with
 152 implications for the fate of ingested carbohydrate.

The primary sources of carbohydrate supporting exercise metabolism are muscle glycogen, and circulating glucose and lactate⁽³⁴⁾. In the fasted state, almost all the circulating glucose is derived from hepatic glycogenolysis and gluconeogenesis, with minor contributions from the kidneys and intestine⁽⁵⁾. When compared to the capacity to store fat, the relatively limited capacity for humans to store carbohydrate has implications for the ability to sustain moderate-to-high intensity exercise⁽⁵⁾. Even amongst lean individuals (~10% body fat), sufficient energy is stored as fat in adipose tissue to theoretically sustain moderate-to-high intensity exercise for many weeks. However, utilising fat as a fuel has a number of limitations in the context of exercise performance. Fat is a relatively “slow” fuel; the rate of adenosine triphosphate (ATP) resynthesis with fat is at least half that when utilising muscle glycogen^(42; 43). Fat is also an “inefficient” fuel on an oxygen basis, requiring ~10% more oxygen consumption for an equivalent energy yield as glucose⁽⁴⁴⁾. Consequently, during high-intensity exercise where rapid ATP resynthesis is required and/or muscle oxygen availability could be limiting, there are advantages to oxidising carbohydrates over fats. Finally, recent evidence implies that glycogen is more than just a fuel and is an important signalling molecule⁽⁴⁵⁾. Low glycogen concentrations in the intramyofibrillar region are associated with impaired sarcoplasmic reticulum calcium release rates and excitation contraction coupling⁽⁴⁶⁾. Therefore, specific depots of glycogen appear to play important roles in both fuelling and regulating skeletal muscle contractile function, hence achieving high carbohydrate availability before and during competition is a goal for athletes competing in almost all endurance sports^(47; 48).

Low carbohydrate (glycogen) availability in muscle and liver is strongly associated with fatigue during prolonged exercise^(49; 50). The amount of glycogen stored in muscle and liver glycogen prior to single, or repeated bouts of exercise positively correlate with subsequent exercise capacity^(49; 50). A number of carbohydrate-related adaptations occur in response to regular endurance training that facilitate improvements in exercise performance. Endurance-trained athletes have a greater capacity to store muscle glycogen, and therefore display an increase in overnight-fasted muscle glycogen concentrations compared to people who are less endurance trained^(5; 51). This increase in basal muscle glycogen concentrations with endurance training is also exaggerated on a high-carbohydrate diet⁽⁵¹⁾, suggesting that endurance-trained athletes can better tolerate high-carbohydrate diets by appropriately storing the excess carbohydrate as muscle glycogen. Interestingly, it seems that basal liver glycogen content does not adapt with endurance training, as endurance-trained athletes tend to exhibit similar liver glycogen concentrations to non-trained controls, when measured in the overnight fasted state⁽⁵⁾. Whether this is also the case in the postprandial state remains to be established.

The increase in fasting muscle (but not liver) glycogen concentrations with endurance training provides trained athletes with a larger depot of glycogen to utilise during exercise and so postpones the point at which critically low muscle glycogen concentrations initiate fatigue. In addition to the

188 greater storage capacity, trained athletes also utilise their muscle glycogen more conservatively
 189 during exercise⁽⁵¹⁾. This sparing of glycogen with endurance training is not specific to muscle, as the
 190 rate of liver glycogen utilisation is also attenuated in endurance-trained athletes compared to controls,
 191 particularly at moderate-to-high exercise intensities⁽⁵⁾. Evidence regarding whether gluconeogenesis
 192 is altered with endurance training is currently equivocal, as some studies indicate endurance training
 193 is associated with an increase in absolute rates of hepatic gluconeogenesis⁽⁵²⁾, whereas others have
 194 shown reductions in hepatic gluconeogenesis after endurance training⁽⁵²⁾. When pooling all the
 195 currently published studies that have concomitantly estimated hepatic glycogenolysis and
 196 gluconeogenesis^(52; 53; 54; 55; 56; 57; 58; 59; 60; 61; 62; 63; 64; 65; 66), it is clear that endurance training is associated
 197 with lower rates of glycogenolysis (**Figure 2A**), whereas any difference in gluconeogenesis with
 198 training status and/or exercise intensity is relatively small and unlikely to be quantitatively important
 199 (**Figure 2B**). Furthermore, it is apparent that hepatic glycogenolysis is the predominant source of
 200 blood glucose during exercise in an overnight fasted state, and the increase in endogenous glucose
 201 appearance with increasing exercise intensity is almost entirely met by an increase in hepatic
 202 glycogenolysis, rather than gluconeogenesis (**Figure 2**).
 203




Figure 2. Hepatic glycogenolysis (**A**) and gluconeogenesis (**B**) in endurance trained individuals and untrained individuals. Each dot represents a group of participants or exercise intensity from a study, and error bars represent 95% confidence intervals (only calculated when published data were available to permit this). The shaded areas represent the 95% confidence intervals of the trend lines. Data are from references: 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66.

204 Interactions between carbohydrate ingestion and exercise occur on multiple levels and in both
 205 directions. Ingesting carbohydrates during exercise can increase total carbohydrate oxidation and
 206 suppress net liver glycogen utilisation and fat oxidation⁽⁶⁷⁾. Whereas even modest exercise potentially
 207 re-directs the metabolic fate of orally ingested sugars. For example, 60 minutes of cycling at 100
 208 watts performed 90 min after fructose ingestion diverts more fructose away from storage (e.g. as
 209 glycogen) and increases fructose oxidation, without altering the conversion of fructose to glucose⁽⁶⁸⁾.

210 This may partly explain why daily exercise can completely prevent the increase in plasma triglyceride
211 concentrations seen with high fructose intakes⁽⁶⁹⁾. Remarkably, the protection offered from exercise
212 against fructose-induced hypertriglyceridemia is seen independently from changes in net energy
213 balance⁽⁶⁹⁾, yet current recommendations for the health effects of dietary sugars rarely consider the
214 context of physical activity status.

215 Since low carbohydrate availability is associated with impaired exercise tolerance, athletes
216 engaging in competitive endurance events regularly consume carbohydrates during exercise⁽⁷⁰⁾. When
217 ingesting glucose alone, the maximal rate at which humans can digest, absorb and metabolise glucose
218 is $\sim 1 \text{ g} \cdot \text{min}^{-1(9)}$, which typically only represents $\sim 44\%$ of total carbohydrate oxidation during exercise
219 at moderate intensity ($\sim 60\% \dot{V}\text{O}_2\text{peak}$) and is therefore insufficient to fully meet the carbohydrate
220 requirements of cycling-based exercise⁽⁷¹⁾. Consequently, oral ingestion of glucose is unable to
221 prevent muscle glycogen depletion during prolonged exercise⁽⁶⁷⁾. It is thought that the primary
222 limitation to the metabolism of orally ingested glucose lies in the splanchnic region, and intestinal
223 absorption of glucose appears to be saturated at $\sim 1 \text{ g} \cdot \text{min}^{-1(9)}$. Ingesting glucose at rates higher than 1
224 $\text{g} \cdot \text{min}^{-1}$ during exercise is therefore likely to lead to accumulation of glucose in the gut and cause
225 gastrointestinal distress. Interestingly, combining fructose with glucose appears to accelerate the
226 digestion, absorption and utilisation of carbohydrate, such that exogenous carbohydrate oxidation
227 rates can reach up to $\sim 1.7 \text{ g} \cdot \text{min}^{-1}$, equating to $\sim 70\%$ of total carbohydrate oxidation^(9; 71). Under these
228 conditions, the relative contribution from endogenous carbohydrate sources is therefore reduced from
229 100% in the fasted state, to $\sim 30\%$ with very high ($2.5 \text{ g} \cdot \text{min}^{-1}$) carbohydrate ingestion rates⁽⁷¹⁾. The
230 primary mechanism by which fructose-glucose mixtures can increase exogenous carbohydrate
231 oxidation over glucose alone is thought to be that intestinal fructose transport utilises a separate
232 pathway than glucose. Specifically, whilst glucose absorption via SGLT-1 is saturated at $\sim 1 \text{ g} \cdot \text{min}^{-1}$,
233 fructose is primarily transported via GLUT5, thereby taking advantage of this alternative pathway
234 and delivering more total carbohydrate to the system⁽⁹⁾.

235

236 **Potential Implications and Applications**

237 ***Exercise Performance***

238 The health and performance implications of carbohydrate intake can be dependent on the specific
239 pathways through which different dietary sugars are absorbed and metabolised. In terms of endurance
240 performance, the accelerated digestion, absorption and utilisation of fructose-glucose mixtures above
241 glucose only, has potential benefits with regards to sparing glycogen stores whilst minimising
242 gastrointestinal distress during exercise⁽⁹⁾. Gastrointestinal complaints are relatively common in
243 endurance events⁽⁷²⁾, which may directly impair performance, but also limit the ability to ingest
244 adequate nutrition to fuel the demands of the exercise. It has recently been demonstrated that the

245 ingestion of either glucose alone, or sucrose (glucose-fructose) can prevent liver glycogen depletion
246 during prolonged (3 h) cycling at a moderate exercise intensity (55% $\text{VO}_{2\text{max}}$)⁽⁶⁷⁾. Whilst there was
247 no further benefit of ingesting sucrose compared to glucose with respect to net liver glycogen
248 depletion, the prevention of liver glycogen depletion with sucrose ingestion was attained with lower
249 ratings of both gut discomfort and perceived exertion, compared to glucose ingestion⁽⁶⁷⁾. Furthermore,
250 when carbohydrates are ingested in large amounts during exercise ($>1.4 \text{ g} \cdot \text{min}^{-1}$), the ingestion of
251 glucose-fructose enhances endurance performance by $\sim 1\text{--}9\%$ more than when glucose is ingested
252 alone⁽⁷³⁾. On the other hand, galactose appears to display relatively low rates of exogenous
253 carbohydrate oxidation during exercise ($\sim 0.4 \text{ g} \cdot \text{min}^{-1}$ oxidised, when ingesting $\sim 1.2 \text{ g} \cdot \text{min}^{-1}$), despite
254 an apparent potential for faster intestinal absorption of galactose compared to glucose in perfusion
255 studies^(74; 75). Moreover, since galactose primarily shares a common intestinal absorption pathway to
256 glucose, combining galactose and glucose ingestion is unlikely to provide the same benefits for
257 exogenous carbohydrate availability and endurance performance as glucose-fructose co-ingestion.

258 In addition to manipulating carbohydrate availability *during* exercise, dietary sugars can also
259 play an important role during post-exercise *recovery*, particularly in multi-stage events such as the
260 Tour de France and the Marathon des Sables, where athletes are required to perform to the best of
261 their ability with less than 24 h of recovery. Under these scenarios, the primary limiting factor to
262 recovery time is glycogen storage rate^(48; 76). Even with high carbohydrate intakes, it is thought to take
263 between 20–24 h to fully restore muscle glycogen concentrations after exhaustive exercise⁽⁷⁷⁾. Thus,
264 on moderate carbohydrate diet, muscle glycogen repletion can take up to 46 h⁽⁷⁸⁾. Therefore, intensive
265 nutritional strategies can be applied to optimise muscle glycogen resynthesis post-exercise. Post-
266 exercise muscle glycogen resynthesis rates display a biphasic response, with the most rapid net
267 synthesis seen within the first 30 min following exercise in an insulin-independent phase⁽⁷⁹⁾.
268 Following this period, muscle glycogen synthesis rates become insulin-dependent and can fall to at
269 least half the rate of that seen within the first 30 min post-exercise⁽⁷⁹⁾. Muscle glycogen resynthesis
270 rates are maximally stimulated with carbohydrate ingestion rates of $\geq 1 \text{ g} \cdot \text{kg body mass}^{-1} \cdot \text{h}^{-1}$ ^(48; 76),
271 and this ingestion rate is also associated with optimal restoration of endurance capacity during short-
272 term (4-h) recovery periods⁽⁷⁶⁾. Therefore, athletes are advised to consume carbohydrate at a rate of
273 $1\text{--}1.2 \text{ g carbohydrate} \cdot \text{kg body mass}^{-1} \cdot \text{h}^{-1}$ during the early stages (4 h) of recovery^(47; 48) and, when
274 these ingestion rates are not achievable, the addition of certain (insulinotropic) proteins, such as milk
275 proteins, to carbohydrate can potentially increase the efficiency of muscle glycogen resynthesis⁽⁸⁰⁾.

276 Current sports nutrition guidelines for recovery do not specify whether the carbohydrate
277 should be from a particular source of sugar (e.g. glucose versus fructose versus galactose)^(47; 48), which
278 is understandable given that muscle glycogen resynthesis rates do not appear to differ whether glucose
279 or glucose-fructose mixtures are ingested^(9; 81), yet overlooks the clear potential for sugars to

differentially affect liver glycogen resynthesis. Indeed, when pooling all currently published data that compare glucose with glucose-fructose mixtures in crossover designs^(50; 81; 82; 83; 84; 85), it is apparent that post-exercise muscle glycogen resynthesis rates do not differ between glucose ingested alone versus glucose-fructose (sucrose) mixtures (**Figure 3A**). Extrapolating these data would suggest that 22 h are required to completely re-synthesise muscle glycogen from a fully depleted state, when following current sports nutrition guidelines, regardless of the type of carbohydrate ingested (**Figure 3A**). This indicates that intestinal absorption of carbohydrate is not rate-limiting to post-exercise muscle glycogen resynthesis. On the other hand, liver glycogen resynthesis appears to be potentially accelerated by glucose-fructose co-ingestion compared to glucose (polymers) alone (**Figure 3B**)^(50; 81; 86), which may be in part due to greater exogenous carbohydrate availability and/or the specific hepatic metabolism of fructose.

A further interesting observation is that liver glycogen resynthesis rates also appear to show a bi-phasic, time-dependent response, albeit in the opposite direction to skeletal muscle. Within the first 2 hours post-exercise, net liver glycogen resynthesis rates are ~30-50% slower than the 3-5 hour period, independent of the type of carbohydrate ingested (2 ± 2 and 5 ± 2 g·h⁻¹ in the 0-2 h post-exercise *versus* 4 ± 2 and 8 ± 2 g·h⁻¹ in the 2-5 h post-exercise, with glucose and sucrose ingestion, respectively)⁽⁸¹⁾. Furthermore, with high rates of carbohydrate ingestion, fructose-glucose mixtures can reduce ratings of gut discomfort during recovery from exercise, compared to glucose ingestion alone⁽⁸¹⁾. Extrapolating these data (i.e. assuming that the first 6.5 h is representative of a full 24-h period) indicates that when only glucose is ingested, complete recovery of liver glycogen stores may take ~25 h (**Figure 3B**). However, when glucose-fructose mixtures are ingested, then liver glycogen repletion could take as little as 11 h (**Figure 3B**). When considering that the time between ending a stage and beginning the next stage in the Tour de France can be ~15 hours, the accelerated recovery of liver glycogen stores with fructose-glucose mixtures is highly meaningful from a practical standpoint.




Figure 3. Studies that have directly compared glucose ingestion alone, with either fructose-glucose or galactose-glucose mixtures, and measure rates of muscle (A) and liver (B) glycogen repletion post-exercise. Each circle represents a timepoint within a study. Error bars represent 95% confidence intervals, and the shaded areas represent the 95% confidence interval of the trend line. For complete recovery of muscle glycogen stores, $600 \text{ mmol} \cdot \text{kgDM}^{-1}$ was chosen on the basis that muscle glycogen concentrations at exhaustion is typically $\sim 115 \text{ mmol} \cdot \text{kgDM}^{-1}$ and the maximal muscle glycogen concentrations of relatively well-trained athletes ($60\text{-}70 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) is between $600\text{-}800 \text{ mmol} \cdot \text{kgDM}^{-1}$ ⁽⁵¹⁾. For complete recovery of liver glycogen stores, 80 g was chosen on the basis that liver glycogen concentrations in the overnight fasted state are $\sim 280 \text{ mmol} \cdot \text{L}^{-1}$. Assuming a liver volume of 1.8 L and the molar mass of a glycosyl unit being 162 g/M , this equates to 80 g glycogen⁽⁵⁾.

311

312

Interestingly, fructose is not the only sugar that more rapidly replenishes liver glycogen contents following exercise than glucose alone. The addition of galactose to glucose also accelerates post-exercise liver glycogen repletion when matched for total carbohydrate intake⁽⁸⁶⁾, and to a similar extent as fructose-glucose ingestion (**Figure 3B**). Since intestinal galactose-glucose absorption should theoretically be slower than fructose-glucose absorption, it is tempting to speculate that the mechanisms by which fructose and galactose enhance liver glycogen resynthesis relate to hepatic metabolism, rather than intestinal absorption. These data also raise the following question: if the Leloir pathway (direct galactose-glucose conversion) is the primary pathway of human galactose metabolism, why is the liver glycogenic response to galactose ingestion more comparable to fructose than to glucose? With regards to generating useful data for applied practice, there is a need to establish the optimal dose and mixture of sugars for rapid liver glycogen resynthesis and whether this translates into improved endurance performance. Accordingly, dose-response studies and direct comparisons of combined galactose-fructose-glucose are warranted.

Whilst the effects of fructose-glucose and galactose-glucose ingestion on glycogen resynthesis are interesting and likely to be important for athletic performance, this will remain speculative in the absence of empirical data. Fortunately, a recent study compared the recovery of exercise capacity with glucose-maltodextrin ingestion versus fructose-maltodextrin ingestion⁽⁸⁷⁾. Since the maltodextrin is hydrolysed, absorbed and oxidised as quickly as free glucose⁽⁹⁾, it can be considered the glucose-maltodextrin is physiologically almost identical to ingesting pure glucose. Athletes were first asked to run on a treadmill at 70% $\dot{V}O_2$ max to exhaustion. Following this, the athletes ingested 90 g carbohydrate per hour during a 4-hour recovery period as either a glucose-maltodextrin mixture, or fructose-maltodextrin mixture. After the recovery period, the athletes ran on the treadmill again at 70% $\dot{V}O_2$ max to exhaustion. During the glucose-maltodextrin trial, the second-bout capacity for these athletes to run was 61.4 ± 9.6 minutes, whereas, when fructose-maltodextrin was ingested in the recovery period, these athletes ran for 81.4 ± 22.3 minutes representing an improvement in second-bout endurance capacity of $\sim 30\%$ ⁽⁸⁷⁾. This provides the first evidence that fructose-glucose ingestion accelerates recovery of exercise capacity. When considered in light of the consistently reported acceleration of liver glycogen recovery, it may be sensible for athletes requiring rapid recovery during multi-stage events to consume fructose-glucose mixtures rather than glucose only. In terms of applying this in practice, it could mean the use of fruit smoothies to supplement carbohydrate intake rather than the commonly held view that pasta is a preferable choice for carbohydrate loading.

Metabolic Health

347 The impact of dietary sugars on hepatic metabolism also has potential metabolic health
348 consequences. Public health recommendations to limit intake of free sugars are primarily based on
349 the effects of diets high in free sugars on body weight and associations with dental caries⁽¹¹⁾. However,
350 distinct metabolic effects of fructose in particular receive much interest with respect to metabolic
351 health. Metabolic health is typically characterised by the ability to maintain relatively stable blood
352 glucose and lipid concentrations in the postprandial state⁽⁸⁸⁾, since high postprandial glucose and/or
353 triglycerides concentrations are associated with cardiovascular disease^(89; 90). The ability to maintain
354 relatively stable circulating metabolite concentrations with relatively little need for insulin represents
355 an important aspect of insulin sensitivity, which is thought to be a fundamental mechanism by which
356 metabolic health is sustained. Whilst insulin sensitivity is most commonly associated with blood
357 glucose control, the many regulatory roles of insulin mean that insulin sensitivity is best considered
358 with respect to the tissue of interest and function of interest. For example, insulin sensitivity of
359 skeletal muscle to glucose uptake, insulin sensitivity of the liver to glucose output, or insulin
360 sensitivity of adipose tissue to lipolysis, etc. This is relevant when discussing the role of fructose in
361 metabolic health as it is apparent that most of the metabolic effects of fructose occur in a tissue-
362 specific manner.

363 The addition of fructose to other ingested nutrients can impact both postprandial glucose and
364 lipid metabolism. Low doses of fructose can in fact lower postprandial glycaemia via increased
365 hepatic glucose disposal secondary to fructose-1-phosphate antagonisation of glucokinase regulatory
366 protein, and thereby enhanced hepatic glucokinase activity^(91; 92; 93). However, compared to glucose
367 ingestion, fructose can enhance postprandial triglyceride concentrations acutely⁽¹⁹⁾ and
368 supplementation of fructose over days/weeks can increase fasting plasma glucose, insulin and
369 triglyceride concentrations, and increase liver fat content, particularly in overweight/obese
370 populations and during positive energy balance^(94; 95). However, some have shown that a positive
371 energy balance and/or saturated fat intake are more potent drivers of liver fat accumulation than
372 specific effects of fructose over glucose^(96; 97). The mechanisms underlying these metabolic changes
373 with fructose ingestion, are thought to include a suppression of hepatic insulin sensitivity to glucose
374 output, stimulation of *de novo* lipogenesis via activation of pyruvate dehydrogenase (upregulated
375 when glycogen concentrations are high⁽⁹⁸⁾), and a reduction in hepatic fatty acid oxidation, leading to
376 increased net lipid synthesis and VLDL-triglyceride production and secretion^(10; 69; 95; 99). This is
377 consistent with data pertaining to post-exercise glycogen resynthesis, since it is thought that insulin
378 resistance to skeletal muscle glucose uptake (leading to hyperglycaemia) and *de novo* lipogenesis
379 (leading to hypertriglyceridemia) are upregulated when glycogen stores are saturated^(63; 100; 101).
380 Furthermore, during non-exercise conditions, the increase in postprandial liver glycogen
381 concentrations seen with a 7-day high-glycaemic index diet occurs in tandem with increases in liver

fat content⁽¹⁰²⁾. The proposed relationship between liver glycogen and lipid metabolism supports the idea that regular exercise can obliterate the negative effects of fructose overfeeding in healthy men^(8; 69), since exercise results in rapid glycogen turnover, and there is clear evidence that the carbohydrate deficit from exercise is a key factor in exercise-induced increases in whole-body glucose control⁽¹⁰³⁾.

Whether exercise can be protective against fructose-induced hypertriglyceridaemia and changes in hepatic insulin sensitivity in overweight and obese populations remains to be established. Given the role of glycogen status in metabolic health, it could be speculated that, when metabolic control is the primary aim, the avoidance of carbohydrates (and in particular fructose-containing sugars) for periods before, during, and/or after exercise could better maintain some of the insulin-sensitising effects of exercise via greater liver glycogen depletion and delayed liver glycogen repletion (**Figure 3**), but this has never been empirically assessed. Fructose can therefore induce changes that are associated with impaired metabolic health, but this appears to be primarily in sedentary, overweight and obese individuals, and when in a positive energy balance. There is evidence that regular exercise has the potential to protect against most (if not all) of these metabolic effects, at least in healthy men. Research is required to determine whether exercise can be protective against metabolic changes with fructose supplementation in people at risk of metabolic disease, and if so, then to characterise the lowest “dose” of exercise that is protective.

Summary and Conclusions

The liver is a primary site of carbohydrate metabolism and particularly the metabolism of fructose and galactose-containing sugars. Hepatic metabolism plays a key role in metabolic health and endurance exercise performance, by assisting in the maintenance of blood glucose and lipid homeostasis during rapid changes in the supply and demand for energy, such as with fasting-feeding cycles and with physical activity. In the fasted state, the liver provides almost all the glucose necessary to maintain blood glucose concentrations during exercise. As exercise intensity increases, thereby accelerating the demand for glucose by skeletal muscle, the increase in liver glucose output is primarily met by releasing stored glucose from glycogen, rather than by increases in *de novo* synthesis of glucose by gluconeogenesis. Similarly, the reduction in liver glucose output during exercise seen in endurance-trained athletes compared to untrained controls, is primarily driven by a reduction in glycogenolysis, as opposed to changes in gluconeogenesis. Therefore, prolonged exercise of a moderate-to-high intensity leads to a depletion of liver glycogen stores unless carbohydrate is ingested during exercise, particularly in less-trained individuals.

For endurance athletes who require rapid recovery for subsequent competitive events, restoration of skeletal muscle and liver glycogen stores are a primary goal. Carbohydrate ingestion is a requirement to replenish glycogen stores within a 24-h timeframe, and ingesting carbohydrate at a

417 rate of $\sim 1 \text{ g}\cdot\text{kg body mass}\cdot\text{h}^{-1}$ within the early (0-4 h) recovery period can assist in optimising this
418 process. Whilst muscle glycogen repletion appears to be largely unaffected by the specific presence
419 of fructose in ingested carbohydrates, liver glycogen repletion rates are potentially enhanced by the
420 ingestion of fructose- or galactose-containing sugars, when compared to glucose alone. There is
421 evidence that the complete restoration of liver glycogen stores after exhaustive exercise could be
422 accelerated by as much as two-fold with the ingestion of fructose-glucose mixtures, compared to
423 glucose-only carbohydrates. Therefore, athletes with multiple competitive events within a 24-h period
424 should aim to consume $\sim 1 \text{ g}\cdot\text{kg body mass}\cdot\text{h}^{-1}$ of carbohydrate with foods providing fructose and
425 glucose. Not only does this enhance restoration of liver (and therefore total body) glycogen stores,
426 there is now evidence that this can reduce the gut discomfort associated with high-carbohydrate
427 ingestion rates, and improve endurance running capacity. There is, however further work required to
428 establish the optimal dose and mixture of carbohydrates to be ingested to maximise post-exercise
429 liver glycogen recovery.

430 The rapid restoration of liver glycogen stores is relevant mainly to a small minority of the
431 population engaging in relatively extreme events. Most people are more concerned about their health
432 than competing in an ultra-endurance event. However, the same knowledge gained about the
433 physiological responses to dietary sugars and exercise, particularly hepatic metabolism, can also be
434 applied to improve metabolic health. Fructose containing sugars have been implicated in inducing
435 hyperglycaemia, hypertriglyceridaemia, hepatic insulin resistance and increases in liver fat content,
436 particularly in overweight/obese populations and when in a positive energy balance. Interestingly,
437 there is evidence in young, healthy men that modest amounts of exercise can completely protect
438 against almost all of these potentially deleterious effects of high-fructose intakes, independent of
439 energy balance. The protective effects of exercise may be due to the carbohydrate deficit and/or
440 glycogen turnover in the liver and skeletal muscle induced by physical activity. Accordingly,
441 specifically avoiding carbohydrates at key times: either before, during and/or after exercise to
442 augment and preserve a glycogen deficit, could be a strategy to enhance metabolic health. However,
443 it is not known if exercise can be protective in populations at risk of metabolic disease, which should
444 be a future research priority.

445 **Acknowledgments.**

446 None

447 **Financial Support**

448 J. T. Gonzalez has received research funding from The European Society of Clinical Nutrition and
449 Metabolism, The Rank Prize Funds, The Physiological Society (UK), The Biotechnology and
450 Biological Sciences Research Council (UK), The Medical Research Council (UK), Arla Foods
451 Ingredients, Lucozade Ribena Suntory and Kenniscentrum Suiker and Voeding, and has acted as a
452 consultant to PepsiCo.

453 J. A. Betts has received research funding from GlaxoSmithKline Nutritional Healthcare R&D,
454 Lucozade Ribena Suntory and Kelloggs, has acted as a consultant for Lucozade Ribena Suntory and
455 PepsiCo and is a scientific advisor to the International Life Sciences Institute (ILSI) Europe Task
456 Force on Energy Balance.

457 This research received no specific grant from any funding agency, commercial or not-for-profit
458 sectors.

459 **Conflict of Interest**

460 None.

461

462 **Authorship**

463 J. T. Gonzalez wrote the initial draft of the manuscript. All authors read, edited and approved of the
464 final manuscript.

465

466

467 **References**

- 468 1. Daston L (2005) Scientific error and the ethos of belief. *Social Research* **72**, 18.
- 469 2. Young FG (1957) Claude Bernard and the discovery of glycogen: A century of retrospect. *Br Med*
- 470 *J* **1**, 1431-1437.
- 471 3. Bernard C (1855) Leçons de physiologie expérimentale appliquée à la médecine, 1: faites au
- 472 Collège de France. *Chez J-B Baillière*.
- 473 4. Bernard C (1879) Leçons sur les phénomènes de la vie communs aux animaux et aux végétaux,
- 474 vol. 2. Baillière, Paris.
- 475 5. Gonzalez JT, Fuchs CJ, Betts JA *et al.* (2016) Liver glycogen metabolism during and after
- 476 prolonged endurance-type exercise. *Am J Physiol Endocrinol Metab* **311**, E543-553.
- 477 6. Jang C, Hui S, Lu W *et al.* (2018) The Small Intestine Converts Dietary Fructose into Glucose and
- 478 Organic Acids. *Cell Metab* **27**, 351-361.e353.
- 479 7. Mutel E, Gautier-Stein A, Abdul-Wahed A *et al.* (2011) Control of blood glucose in the absence
- 480 of hepatic glucose production during prolonged fasting in mice: induction of renal and intestinal
- 481 gluconeogenesis by glucagon. *Diabetes* **60**, 3121-3131.
- 482 8. Gonzalez JT, Betts JA (2018) Dietary Fructose Metabolism By Splanchnic Organs: Size Matters.
- 483 *Cell Metab* **27**, 483-485.
- 484 9. Gonzalez JT, Fuchs CJ, Betts JA *et al.* (2017) Glucose Plus Fructose Ingestion for Post-Exercise
- 485 Recovery-Greater than the Sum of Its Parts? *Nutrients* **9**, E344-359.
- 486 10. Tappy L, Lê KA (2010) Metabolic effects of fructose and the worldwide increase in obesity.
- 487 *Physiol Rev* **90**, 23-46.
- 488 11. World Health Organization (2015) Guideline: Sugars intake for adults and children. Geneva.
- 489 12. Scientific Advisory Committee on Nutrition (2015) Carbohydrates and Health. London: TSO.
- 490 13. Hawley JA, Dennis SC, Laidler BJ *et al.* (1991) High rates of exogenous carbohydrate oxidation
- 491 from starch ingested during prolonged exercise. *J Appl Physiol* (1985) **71**, 1801-1806.
- 492 14. Daniel H, Zietek T (2015) Taste and move: glucose and peptide transporters in the gastrointestinal
- 493 tract. *Exp Physiol* **100**, 1441-1450.
- 494 15. Kelley D, Mitrakou A, Marsh H *et al.* (1988) Skeletal muscle glycolysis, oxidation, and storage
- 495 of an oral glucose load. *J Clin Invest* **81**, 1563-1571.
- 496 16. Ferrannini E, Bjorkman O, Reichard GA *et al.* (1985) The disposal of an oral glucose load in
- 497 healthy subjects. A quantitative study. *Diabetes* **34**, 580-588.
- 498 17. Williams CA (1986) Metabolism of lactose and galactose in man. *Prog Biochem Pharmacol* **21**,
- 499 219-247.
- 500 18. Ercan N, Nuttall FQ, Gannon MC *et al.* (1993) Effects of glucose, galactose, and lactose ingestion
- 501 on the plasma glucose and insulin response in persons with non-insulin-dependent diabetes mellitus.
- 502 *Metabolism* **42**, 1560-1567.
- 503 19. Chong MF, Fielding BA, Frayn KN (2007) Mechanisms for the acute effect of fructose on
- 504 postprandial lipemia. *Am J Clin Nutr* **85**, 1511-1520.
- 505 20. Heinz F, Lamprecht W, Kirsch J (1968) Enzymes of fructose metabolism in human liver. *J Clin*
- 506 *Invest* **47**, 1826-1832.
- 507 21. Adelman RC, Ballard FJ, Weinhouse S (1967) Purification and properties of rat liver fructokinase.
- 508 *J Biol Chem* **242**, 3360-3365.
- 509 22. Hers HG, Kusaka T (1953) The metabolism of fructose-1-phosphate in the liver. *Biochim Biophys*
- 510 *Acta* **11**, 427-437.
- 511 23. Mayes PA (1993) Intermediary metabolism of fructose. *Am J Clin Nutr* **58**, 754S-765S.
- 512 24. Björkman O, Gunnarsson R, Hagström E *et al.* (1989) Splanchnic and renal exchange of infused
- 513 fructose in insulin-deficient type 1 diabetic patients and healthy controls. *J Clin Invest* **83**, 52-59.
- 514 25. Bar-On H, Stein Y (1968) Effect of glucose and fructose administration on lipid metabolism in
- 515 the rat. *J Nutr* **94**, 95-105.
- 516 26. Carmona A, Freedland RA (1989) Comparison among the lipogenic potential of various
- 517 substrates in rat hepatocytes: the differential effects of fructose-containing diets on hepatic
- 518 lipogenesis. *J Nutr* **119**, 1304-1310.

- 519 27. Sanders FWB, Acharjee A, Walker C *et al.* (2018) Hepatic steatosis risk is partly driven by
520 increased de novo lipogenesis following carbohydrate consumption. *Genome Biol* **19**, 79.
- 521 28. Timson DJ, Reece RJ (2003) Functional analysis of disease-causing mutations in human
522 galactokinase. *Eur J Biochem* **270**, 1767-1774.
- 523 29. Sørensen M, Mikkelsen KS, Frisch K *et al.* (2011) Hepatic galactose metabolism quantified in
524 humans using 2-18F-fluoro-2-deoxy-D-galactose PET/CT. *J Nucl Med* **52**, 1566-1572.
- 525 30. Sunehag AL, Haymond MW (2002) Splanchnic galactose extraction is regulated by coingestion
526 of glucose in humans. *Metabolism* **51**, 827-832.
- 527 31. Coss-Bu JA, Sunehag AL, Haymond MW (2009) Contribution of galactose and fructose to
528 glucose homeostasis. *Metabolism* **58**, 1050-1058.
- 529 32. Niewoehner CB, Neil B, Martin T (1990) Hepatic uptake and metabolism of oral galactose in
530 adult fasted rats. *Am J Physiol* **259**, E804-813.
- 531 33. Décombaz J, Jentjens R, Ith M *et al.* (2011) Fructose and galactose enhance postexercise human
532 liver glycogen synthesis. *Med Sci Sports Exerc* **43**, 1964-1971.
- 533 34. van Loon LJ, Greenhaff PL, Constantin-Teodosiu D *et al.* (2001) The effects of increasing
534 exercise intensity on muscle fuel utilisation in humans. *J Physiol* **536**, 295-304.
- 535 35. Achten J, Venables MC, Jeukendrup AE (2003) Fat oxidation rates are higher during running
536 compared with cycling over a wide range of intensities. *Metabolism* **52**, 747-752.
- 537 36. Chen YC, Travers RL, Walhin JP *et al.* (2017) Feeding influences adipose tissue responses to
538 exercise in overweight men. *Am J Physiol Endocrinol Metab* **313**, E84-E93.
- 539 37. Gonzalez JT, Veasey RC, Rumbold PL *et al.* (2013) Breakfast and exercise contingently affect
540 postprandial metabolism and energy balance in physically active males. *Br J Nutr* **110**, 721-732.
- 541 38. Edinburgh RM, Hengist A, Smith HA *et al.* (2018) Pre-exercise breakfast ingestion versus
542 extended overnight fasting increases postprandial glucose flux after exercise in healthy men.
543 *American Journal of Physiology – Endocrinology and Metabolism* **Accepted-ahead-of-print** doi:
544 10.1152/ajpendo.00163.2018.
- 545 39. van Loon LJ, Jeukendrup AE, Saris WH *et al.* (1999) Effect of training status on fuel selection
546 during submaximal exercise with glucose ingestion. *J Appl Physiol* (1985) **87**, 1413-1420.
- 547 40. Fletcher G, Eves FF, Glover EI *et al.* (2017) Dietary intake is independently associated with the
548 maximal capacity for fat oxidation during exercise. *Am J Clin Nutr* **105**, 864-872.
- 549 41. Romijn JA, Coyle EF, Sidossis LS *et al.* (1993) Regulation of endogenous fat and carbohydrate
550 metabolism in relation to exercise intensity and duration. *Am J Physiol* **265**, E380-391.
- 551 42. Walter G, Vandenborne K, Elliott M *et al.* (1999) In vivo ATP synthesis rates in single human
552 muscles during high intensity exercise. *J Physiol* **519 Pt 3**, 901-910.
- 553 43. Hultman E, Harris RC (1988) *Carbohydrate metabolism, Principles of Exercise Biochemistry*.
554 Basel, Switzerland: S.Karger.
- 555 44. Jeukendrup AE, Wallis GA (2005) Measurement of substrate oxidation during exercise by means
556 of gas exchange measurements. *Int J Sports Med* **26 Suppl 1**, S28-37.
- 557 45. Ørtenblad N, Westerblad H, Nielsen J (2013) Muscle glycogen stores and fatigue. *J Physiol* **591**,
558 4405-4413.
- 559 46. Ørtenblad N, Nielsen J (2015) Muscle glycogen and cell function--Location, location, location.
560 *Scand J Med Sci Sports* **25 Suppl 4**, 34-40.
- 561 47. Thomas DT, Erdman KA, Burke LM (2016) American College of Sports Medicine Joint Position
562 Statement. Nutrition and Athletic Performance. *Med Sci Sports Exerc* **48**, 543-568.
- 563 48. Burke LM, van Loon LJC, Hawley JA (2017) Postexercise muscle glycogen resynthesis in
564 humans. *J Appl Physiol* (1985) **122**, 1055-1067.
- 565 49. Bergström J, Hermansen L, Hultman E *et al.* (1967) Diet, muscle glycogen and physical
566 performance. *Acta Physiol Scand* **71**, 140-150.
- 567 50. Casey A, Mann R, Banister K *et al.* (2000) Effect of carbohydrate ingestion on glycogen
568 resynthesis in human liver and skeletal muscle, measured by (13)C MRS. *Am J Physiol Endocrinol*
569 *Metab* **278**, E65-75.

570 51. Areta JL, Hopkins WG (2018) Skeletal Muscle Glycogen Content at Rest and During Endurance
571 Exercise in Humans: A Meta-Analysis. *Sports Med*, **48**, 2091-2102.

572 52. Bergman BC, Horning MA, Casazza GA *et al.* (2000) Endurance training increases
573 gluconeogenesis during rest and exercise in men. *Am J Physiol Endocrinol Metab* **278**, E244-251.

574 53. Wahren J, Felig P, Ahlborg G *et al.* (1971) Glucose metabolism during leg exercise in man. *J*
575 *Clin Invest* **50**, 2715-2725.

576 54. Ahlborg G, Felig P, Hagenfeldt L *et al.* (1974) Substrate turnover during prolonged exercise in
577 man. Splanchnic and leg metabolism of glucose, free fatty acids, and amino acids. *J Clin Invest* **53**,
578 1080-1090.

579 55. Sestoft L, Trap-Jensen J, Lyngsooe J *et al.* (1977) Regulation of gluconeogenesis and ketogenesis
580 during rest and exercise in diabetic subjects and normal men. *Clin Sci Mol Med* **53**, 411-418.

581 56. Stanley WC, Wisneski JA, Gertz EW *et al.* (1988) Glucose and lactate interrelations during
582 moderate-intensity exercise in humans. *Metabolism* **37**, 850-858.

583 57. Friedlander AL, Casazza GA, Horning MA *et al.* (1997) Training-induced alterations of glucose
584 flux in men. *J Appl Physiol* (1985) **82**, 1360-1369.

585 58. Friedlander AL, Casazza GA, Horning MA *et al.* (1998) Training-induced alterations of
586 carbohydrate metabolism in women: women respond differently from men. *J Appl Physiol* (1985) **85**,
587 1175-1186.

588 59. Lavoie C, Ducros F, Bourque J *et al.* (1997) Glucose metabolism during exercise in man: the role
589 of insulin and glucagon in the regulation of hepatic glucose production and gluconeogenesis. *Can J*
590 *Physiol Pharmacol* **75**, 26-35.

591 60. Trimmer JK, Schwarz JM, Casazza GA *et al.* (2002) Measurement of gluconeogenesis in
592 exercising men by mass isotopomer distribution analysis. *J Appl Physiol* (1985) **93**, 233-241.

593 61. Roef MJ, Kalhan SC, Reijngoud DJ *et al.* (2002) Lactate disposal via gluconeogenesis is increased
594 during exercise in patients with mitochondrial myopathy due to complex I deficiency. *Pediatr Res*
595 **51**, 592-597.

596 62. Roef MJ, de Meer K, Kalhan SC *et al.* (2003) Gluconeogenesis in humans with induced
597 hyperlactatemia during low-intensity exercise. *Am J Physiol Endocrinol Metab* **284**, E1162-1171.

598 63. Petersen KF, Price TB, Bergeron R (2004) Regulation of net hepatic glycogenolysis and
599 gluconeogenesis during exercise: impact of type 1 diabetes. *J Clin Endocrinol Metab* **89**, 4656-4664.

600 64. Huidekoper HH, Ackermans MT, Koopman R *et al.* (2013) Normal rates of whole-body fat
601 oxidation and gluconeogenesis after overnight fasting and moderate-intensity exercise in patients
602 with medium-chain acyl-CoA dehydrogenase deficiency. *J Inherit Metab Dis* **36**, 831-840.

603 65. Emhoff CA, Messonnier LA, Horning MA *et al.* (2013) Gluconeogenesis and hepatic
604 glycogenolysis during exercise at the lactate threshold. *J Appl Physiol* (1985) **114**, 297-306.

605 66. Webster CC, Noakes TD, Chacko SK *et al.* (2016) Gluconeogenesis during endurance exercise
606 in cyclists habituated to a long-term low carbohydrate high fat diet. *J Physiol* **594**, 4389-4405.

607 67. Gonzalez JT, Fuchs CJ, Smith FE *et al.* (2015) Ingestion of glucose or sucrose prevents liver but
608 not muscle glycogen depletion during prolonged endurance-type exercise in trained cyclists. *Am J*
609 *Physiol Endocrinol Metab* **309**, E1032-1039.

610 68. Egli L, Lecoultre V, Cros J *et al.* (2016) Exercise performed immediately after fructose ingestion
611 enhances fructose oxidation and suppresses fructose storage. *Am J Clin Nutr* **103**, 348-355.

612 69. Egli L, Lecoultre V, Theytaz F *et al.* (2013) Exercise prevents fructose-induced
613 hypertriglyceridemia in healthy young subjects. *Diabetes* **62**, 2259-2265.

614 70. Saris WH, van Erp-Baart MA, Brouns F *et al.* (1989) Study on food intake and energy expenditure
615 during extreme sustained exercise: the Tour de France. *Int J Sports Med* **10 Suppl 1**, S26-31.

616 71. Jentjens RL, Achten J, Jeukendrup AE (2004) High oxidation rates from combined carbohydrates
617 ingested during exercise. *Med Sci Sports Exerc* **36**, 1551-1558.

618 72. Pfeiffer B, Stellingwerff T, Hodgson AB *et al.* (2012) Nutritional intake and gastrointestinal
619 problems during competitive endurance events. *Med Sci Sports Exerc* **44**, 344-351.

620 73. Rowlands DS, Houltham S, Musa-Veloso K *et al.* (2015) Fructose-Glucose Composite
621 Carbohydrates and Endurance Performance: Critical Review and Future Perspectives. *Sports Med* **45**,
622 1561-1576.

623 74. Holdsworth CD, Dawson AM (1964) The absorption of monosaccharides in man. *Clin Sci* **27**,
624 371-379.

625 75. Cook GC (1977) Rates and mechanisms of glucose, galactose, and xylose absorption in man in
626 vivo. *Scand J Gastroenterol* **12**, 733-737.

627 76. Alghannam AF, Gonzalez JT, Betts JA (2018) Restoration of Muscle Glycogen and Functional
628 Capacity: Role of Post-Exercise Carbohydrate and Protein Co-Ingestion. *Nutrients* **10**, E253-280.

629 77. Coyle EF (1991) Timing and method of increased carbohydrate intake to cope with heavy
630 training, competition and recovery. *J Sports Sci* **9 Spec No**, 29-51; discussion 51-22.

631 78. Piehl K (1974) Time course for refilling of glycogen stores in human muscle fibres following
632 exercise-induced glycogen depletion. *Acta Physiol Scand* **90**, 297-302.

633 79. Price TB, Rothman DL, Taylor R *et al.* (1994) Human muscle glycogen resynthesis after exercise:
634 insulin-dependent and -independent phases. *J Appl Physiol* (1985) **76**, 104-111.

635 80. Betts JA, Williams C (2010) Short-term recovery from prolonged exercise: exploring the potential
636 for protein ingestion to accentuate the benefits of carbohydrate supplements. *Sports Med* **40**, 941-
637 959.

638 81. Fuchs CJ, Gonzalez JT, Beelen M *et al.* (2016) Sucrose ingestion after exhaustive exercise
639 accelerates liver, but not muscle glycogen repletion compared with glucose ingestion in trained
640 athletes. *J Appl Physiol* (1985) **120**, 1328-1334.

641 82. Blom PC, Høstmark AT, Vaage O *et al.* (1987) Effect of different post-exercise sugar diets on
642 the rate of muscle glycogen synthesis. *Med Sci Sports Exerc* **19**, 491-496.

643 83. Bowtell JL, Gelly K, Jackman ML *et al.* (2000) Effect of different carbohydrate drinks on whole
644 body carbohydrate storage after exhaustive exercise. *J Appl Physiol* (1985) **88**, 1529-1536.

645 84. Wallis GA, Hulston CJ, Mann CH *et al.* (2008) Postexercise muscle glycogen synthesis with
646 combined glucose and fructose ingestion. *Med Sci Sports Exerc* **40**, 1789-1794.

647 85. Trommelen J, Beelen M, Pinckaers PJ *et al.* (2016) Fructose Coingestion Does Not Accelerate
648 Postexercise Muscle Glycogen Repletion. *Med Sci Sports Exerc* **48**, 907-912.

649 86. Decombaz J, Jentjens R, Ith M *et al.* (2011) Fructose and galactose enhance postexercise human
650 liver glycogen synthesis. *Med Sci Sports Exerc* **43**, 1964-1971.

651 87. Maunder E, Podlogar T, Wallis GA (2018) Postexercise Fructose-Maltodextrin Ingestion
652 Enhances Subsequent Endurance Capacity. *Med Sci Sports Exerc* **50**, 1039-1045.

653 88. Edinburgh RM, Betts JA, Burns SF *et al.* (2017) Concordant and divergent strategies to improve
654 postprandial glucose and lipid metabolism. *Nutrition Bulletin* **42**, 113-122.

655 89. Nordestgaard BG, Benn M, Schnohr P *et al.* (2007) Nonfasting triglycerides and risk of
656 myocardial infarction, ischemic heart disease, and death in men and women. *J Am Med Assoc* **298**,
657 299-308.

658 90. Ning F, Zhang L, Dekker JM *et al.* (2012) Development of coronary heart disease and ischemic
659 stroke in relation to fasting and 2-hour plasma glucose levels in the normal range. *Cardiovasc*
660 *Diabetol* **11**, 76.

661 91. Van Schaftingen E, Detheux M, Veiga da Cunha M (1994) Short-term control of glucokinase
662 activity: role of a regulatory protein. *FASEB J* **8**, 414-419.

663 92. Donmoyer CM, Ejiofor J, Lacy DB *et al.* (2001) Fructose augments infection-impaired net hepatic
664 glucose uptake during TPN administration. *Am J Physiol Endocrinol Metab* **280**, E703-711.

665 93. Sievenpiper JL, Chiavaroli L, de Souza RJ *et al.* (2012) 'Catalytic' doses of fructose may benefit
666 glycaemic control without harming cardiometabolic risk factors: a small meta-analysis of randomised
667 controlled feeding trials. *Br J Nutr* **108**, 418-423.

668 94. Stanhope KL, Schwarz JM, Keim NL *et al.* (2009) Consuming fructose-sweetened, not glucose-
669 sweetened, beverages increases visceral adiposity and lipids and decreases insulin sensitivity in
670 overweight/obese humans. *J Clin Invest* **119**, 1322-1334.

671 95. Lecoultre V, Egli L, Carrel G *et al.* (2013) Effects of fructose and glucose overfeeding on hepatic
 672 insulin sensitivity and intrahepatic lipids in healthy humans. *Obesity (Silver Spring)* **21**, 782-785.
 673 96. Johnston RD, Stephenson MC, Crossland H *et al.* (2013) No difference between high-fructose
 674 and high-glucose diets on liver triacylglycerol or biochemistry in healthy overweight men.
 675 *Gastroenterology* **145**, 1016-1025.e1012.
 676 97. Luukkonen PK, Sädevirta S, Zhou Y *et al.* (2018) Saturated Fat Is More Metabolically Harmful
 677 for the Human Liver Than Unsaturated Fat or Simple Sugars. *Diabetes Care* **41**, 1732-1739.
 678 98. Kiilerich K, Gudmundsson M, Birk JB *et al.* (2010) Low muscle glycogen and elevated plasma
 679 free fatty acid modify but do not prevent exercise-induced PDH activation in human skeletal muscle.
 680 *Diabetes* **59**, 26-32.
 681 99. Park OJ, Cesar D, Faix D *et al.* (1992) Mechanisms of fructose-induced hypertriglyceridaemia in
 682 the rat. Activation of hepatic pyruvate dehydrogenase through inhibition of pyruvate dehydrogenase
 683 kinase. *Biochem J* **282** (Pt 3), 753-757.
 684 100. Rabøl R, Petersen KF, Dufour S *et al.* (2011) Reversal of muscle insulin resistance with exercise
 685 reduces postprandial hepatic de novo lipogenesis in insulin resistant individuals. *Proc Natl Acad Sci*
 686 *U S A* **108**, 13705-13709.
 687 101. Nolte LA, Gulve EA, Holloszy JO (1994) Epinephrine-induced in vivo muscle glycogen
 688 depletion enhances insulin sensitivity of glucose transport. *J Appl Physiol (1985)* **76**, 2054-2058.
 689 102. Bawden S, Stephenson M, Falcone Y *et al.* (2017) Increased liver fat and glycogen stores after
 690 consumption of high versus low glycaemic index food: A randomized crossover study. *Diabetes Obes*
 691 *Metab* **19**, 70-77.
 692 103. Taylor HL, Wu CL, Chen YC *et al.* (2018) Post-Exercise Carbohydrate-Energy Replacement
 693 Attenuates Insulin Sensitivity and Glucose Tolerance the Following Morning in Healthy Adults.
 694 *Nutrients* **10**, E123-134.
 695
 696